

Contents lists available at ScienceDirect

Science of the Total Environment



Endangered Australian top predator is frequently exposed to anticoagulant rodenticides



James M. Pay^{a,*}, Todd E. Katzner^b, Clare E. Hawkins^a, Leon A. Barmuta^a, William E. Brown^c, Jason M. Wiersma^d, Amelia J. Koch^{a,d}, Nick J. Mooney^e, Elissa Z. Cameron^{a,f}

^a School of Natural Sciences, University of Tasmania, Hobart, TAS, Australia

^b US Geological Survey, Forest and Rangeland Ecosystem Science Center, Boise, ID, USA

^c Department of Primary Industries, Parks, Water and Environment, Hobart, TAS, Australia

^d Forest Practices Authority, 30 Patrick St, Hobart, TAS, Australia

^e Birdlife Australia Raptor Group, Birldlife Australia, Carlton, VIC, Australia

^f School of Biological Sciences, University of Canterbury, CHC, New Zealand

HIGHLIGHTS

GRAPHICAL ABSTRACT

- First systematic study of AR exposure in an Australian top predator
- ARs were detected in 74% of 50 eagle carcasses collected between 1996 and 2018.
- AR levels were high in comparison to work on other eagle species.
- AR levels associated with proximity to agriculture and human population density
- ARs may be causing broad contamination of Australia's terrestrial food chains.

ARTICLE INFO

Article history: Received 4 February 2021 Received in revised form 3 May 2021 Accepted 6 May 2021 Available online 10 May 2021

Editor: Damia Barcelo

Keywords: Rodenticide Environmental contamination SGARs Secondary poisoning Predator



ABSTRACT

Anticoagulant rodenticides (ARs) used to control mammalian pest populations cause secondary exposure of predatory species throughout much of the world. It is important to understand the drivers of non-target AR exposure patterns as context for assessing long-term effects and developing effective mitigation for these toxicants. In Australia, however, little is known about exposure and effects of ARs on predators. We detected AR residues in 74% of 50 opportunistically collected carcasses of the Tasmanian wedge-tailed eagle (*Aquila audax fleayi*), an endangered apex predator. In 22% of birds tested, or 31% of those exposed, liver concentrations of second generation ARs (SGARs) were >0.1 mg/kg ww. Eagles were exposed to flocoumafen, a toxicant only available from agricultural suppliers, at an exceptionally high rate (40% of birds tested). Liver SGAR concentrations were positively associated with the proportion of agricultural habitat and human population density in the area around where each eagle died. The high exposure rate in a species not known to regularly prey upon synanthropic rodents supports the hypothesis that apex predators are vulnerable to SGARs. Our results indicate that AR exposure constitutes a previously unrecognized threat to Tasmanian wedge-tailed eagles and highlight the importance of efforts to address non-target AR exposure in Australia.

© 2021 Published by Elsevier B.V.

1. Introduction

* Corresponding author. E-mail address: James.Pay@utas.edu.au (J.M. Pay).

https://doi.org/10.1016/j.scitotenv.2021.147673 0048-9697/© 2021 Published by Elsevier B.V. Anticoagulant rodenticides (ARs) are used worldwide to control mammalian pest populations. These compounds function by inhibiting

blood clotting mechanisms in vertebrates, resulting in internal hemorrhaging (Rattner et al., 2014). The discovery of resistance to the first-generation of ARs (FGARs) led to the development of second-generation ARs (SGARs) in the 1970s (van den Brink et al., 2018). To be lethal, FGARs generally require consecutive intake over several days to accumulate sufficiently high concentrations (Erickson and Urban, 2004). Conversely, SGARs are usually lethal from a single exposure and persist longer in the environment (Erickson and Urban, 2004; van den Brink et al., 2018). The persistence of AR compounds (Horak et al., 2018), the delay in mortality after bait consumption (Lee et al., 2006) and the behavioral changes that occur as a symptom of poisoning (Brakes and Smith, 2005; Mooney, 2017) can make poisoned rodents AR vectors to non-target predatory species.

Detrimental non-target exposure to ARs has been shown in numerous populations of predators in Europe and North America (Christensen et al., 2012; López-Perea et al., 2015; Riley et al., 2007; Shore et al., 2003; Thomas et al., 2017). These effects can be significant, with populationlevel effects from non-target exposure documented for mammals (Jacquot et al., 2013) and raptors (Thomas et al., 2011). It is thought that species that regularly prey upon small rodents are at higher risk of poisoning, due to the likelihood of consuming AR targeted species (Hindmarch and Elliott, 2018). However, the primary consumption of AR baits by non-target species, as well as the potential for SGARs to move through trophic levels, may lead to wider contamination of terrestrial food chains (Hindmarch and Elliott, 2018; Thomas et al., 2011). If such broadscale contamination is apparent, species at higher trophic levels may be at increased risk of AR exposure (Riley et al., 2007; Thomas et al., 2011).

It is necessary to understand the drivers of patterns in non-target AR exposure in order to assess long-term effects and to develop effective mitigation. There are documented differences in AR exposure of predators between the sexes (Mcdonald et al., 1998) and among age groups (Christensen et al., 2012; Ruiz-Suárez et al., 2016). That said, local anthropogenic factors are likely the most significant drivers of overall risk of non-target exposure. For example, human population density and developed surface area have been linked to the probability and level of AR exposure of numerous predators (Lohr, 2018; Lopez-Perea and Mateo, 2018; Nogeire et al., 2015; Serieys et al., 2015). Agricultural AR use has also been suggested as the cause of non-target poisoning of predators (Birks, 1998; Fourel et al., 2018; Hindmarch et al., 2017; Hughes et al., 2013), but only a few recent studies have found empirical evidence of this relationship (Coeurdassier et al., 2019; López-Perea et al., 2018; Rial-Berriel et al., 2021; Sainsbury et al., 2018).

AR use is largely unmonitored in Australia and recent work has highlighted the need for the evaluation of its effects on Australasian taxa (Lohr, 2018; Lohr and Davis, 2018). The Tasmanian wedge-tailed eagle (Aquila audax fleavi) is a subspecies of wedge-tailed eagle endemic to the Australian island of Tasmania (Commonwealth of Australia, 1999). With the loss of the thylacine (Thylacinus cynocephalus) and recent declines in populations of Tasmanian devils (Sarcophilus harrisii), the wedge-tailed eagle serves a particularly important ecological function as one of the few remaining top predators in Tasmanian ecosystems. The subspecies is listed as endangered (Commonwealth of Australia, 1999; State Government of Tasmania, 1995), with conservation concern based upon a series of threats, including nest failures caused by anthropogenic disturbance, low breeding success rates, habitat loss, collisions with anthropogenic structures, lead poisoning, and illegal persecution (Bell and Mooney, 1998; Mooney and Holdsworth, 1991; Pay et al., 2020; Threatened Species Section, 2006).

ARs are not recognized as a significant threat to the Tasmanian wedge-tailed eagle population, as the species generally avoids areas of high human population density, and rodents represent a very small portion of their diet (Marchant and Higgins, 1993). That said, wedge-tailed eagles show a high sensitivity to pindone (Martin et al., 1994), an AR used to control European rabbit (*Oryctolagus cuniculus*) populations, a primary prey species of wedge-tailed eagles (Debus et al., 2007;

Marchant and Higgins, 1993). Furthermore, if ARs are moving through Tasmania's food chains, then the high trophic position of the wedgetailed eagle may increase their susceptibility to exposure to various AR compounds. Finally, because of the long-lived and slow breeding life history strategy of this species, it is likely highly vulnerable to increased mortality rates brought on by toxicants such as ARs.

Our aims in this research were to determine to what extent Tasmanian wedge-tailed eagles are exposed to ARs, and to investigate the factors that influence AR exposure in the population. Specifically, we evaluated (1) liver tissue concentrations of individual ARs known to be used in Tasmania and the total SGAR concentration of each individual eagle; and the relationship between both (2) total liver SGAR concentration and (3) probability of exposure with intrinsic (age and sex) and extrinsic (human population density, agricultural land use, and year of death) factors.

2. Methods

2.1. Study area

This study was conducted on mainland Tasmania, an island state located 240 km south of continental Australia. Tasmania covers an area of 68,150 km², with an estimated human population of 520,830 (Australian Bureau of Statistics, 2018; Fig. 1b). Areas of minimal land use, nature conservation and other protected areas account for 50.3% (34,280 km²) of the Tasmanian land area (DPIPWE, 2015). Agriculture occupies 18,900 km² (27.7%; DPIPWE, 2015) mostly in the north and east of the state (Fig. 1c). The Tasmanian agricultural land area is comprised of modified pastures (75.4%), native vegetation pastures (14.5%), irrigated crops (8.8%), and non-irrigated crops (1%; DPIPWE, 2015).

2.2. Sample collection

Eagles were collected as carcasses found opportunistically throughout Tasmania (Fig. 1a) between 1996 and 2018, by government departments, various industries, and volunteers. All carcasses were placed in -20 °C freezer storage by the Department of Primary Industries, Parks, Water and the Environment (DPIPWE, Threatened Species Section, Hobart, Tasmania) and the Tasmanian Museum and Art Gallery (TMAG, Collection and Research Facility, Rosny, Tasmania). Data recorded for each carcass included location and the date the carcass was found. We thawed the carcasses and harvested tissues from them between May 2017 and March 2018. We collected a whole liver lobe and a muscle sample from each carcass. The tissue samples were stored at -20 °C until sample preparation, when we thawed them at room temperature. We weighed out a 4 g (\pm 0.1 g) wet weight sample from the middle of each liver lobe using a digital balance (precision \pm 0.0001 g (Mettler Toledo, US). New scalpel blades and gloves were used between samples during collection and preparation to prevent cross contamination.

2.3. Residue analysis

2.3.1. Sample preparation

All toxicological analyses were carried out at Edith Cowan University Analytical Facility (Joondalup, Western Australia). Each liver sample was freeze-dried and homogenized. Homogenized samples were transferred into centrifuge plastic tubes (15 ml) and 10 ml of acetonitrile was added to the tubes with a 10 μ l (10 ng/ μ l) solution containing deuterated surrogates. Analytes were extracted using a sonication bath (15 min sonication for each aliquot). After extraction, samples were centrifuged at 3247 relative centrifugal force (rcf) for 5 min, transferred to a new centrifuge tube with 2 ml of hexane, vortexed for 5 min and centrifuged at 3247 rcf for a further 5 min. Each sample was then evaporated and reconstituted in 400 μ l of 50:50 ACN/H20 solution. The final



Fig. 1. a) Location and liver total SGAR concentration threshold for 50 Tasmanian wedge-tailed eagle carcasses collected between 1996 and 2018. Maps b) and c) indicate the spatial distribution of the Tasmanian human population (2016 data; Australian Bureau of Statistics, 2018) and agricultural land use area (2015 data; DPIPWE, 2015) respectively.

extracts were transferred to 2 ml Teflon-lined vials and stored at 0-4 °C until analysis.

2.3.2. LC-MS analysis

Liver samples were analyzed for ARs registered for use in Australia (Australian Pesticides and Veterinary Medicines Authority, 2019). Concentrations of five SGARs (brodifacoum, bromadiolone, difethialone, difenacoum and flocoumafen) and three FGARs (coumatetralyl, pindone and warfarin) (see Appendix Table A.1 for the manufacturers of the analytical standards and surrogates) were evaluated using a TSQ Quantiva triple quadrupole Mass Spectrometer (LC-MS) from Thermo Fisher (Thermo Fisher Scientific Corporation, US) (see Appendix B for details of the chromatographic method). Calibration curves and recovery rates for each analytical run were calculated using organic chicken livers spiked with three working solutions of each analytical standard. Recovery rates for the target ARs averaged 96.75%, whilst limits of detection (LOD) and limits of quantification (LOQ) ranged from 0.0005-0.0125 mg/kg and 0.001-0.025 mg/kg respectively (Appendix Table A.2). All detections that were >LOD but <LOQ were reported as present at trace levels. Three organic chicken liver blanks were included in each run to monitor cross-contamination. Every 10th sample was reinjected for a duplicate read (average percentage relative standard deviation of recoveries (RSD) 4.1%) and duplicate blind sample extractions were carried out for five randomly selected samples (average RSD 4.1%). Concentrations were reported on a dry weight basis (mg/kg dw).

2.4. Potential drivers of AR exposure

We evaluated potential drivers of AR exposure as a response to a suite of intrinsic and extrinsic explanatory variables. The intrinsic variables we considered were the sex of the bird (determined genetically using muscle tissue; Appendix C) and its age (broadly characterized into adults and pre-adults based on plumage; Appendix D). Extrinsic explanatory variables were the year in which the carcass was found, and both the mean human population density per km² (Australian Bureau of Statistics, 2018) and the proportion of agricultural area (DPIPWE, 2015) in the area surrounding where each carcass was found. Areas we categorized as agricultural included all types of animal production (intensive animal production, native vegetation grazing, and modified pastures grazing) and all types of horticulture (both non-irrigated, and irrigated cropping; see Appendix E). We defined the area around where each carcass was found based on the size of the estimated home range of adult and pre-adult eagles (25 km² for adults and 420 km² for pre-adults; see Appendix F). We buffered each carcass location by an area corresponding to the age-specific home range and calculated the mean human population density per km² and the proportion of agricultural land within the buffered area. To maximize accuracy in estimates of spatial predictor variables, both human population density and agricultural land use area were calculated from data as close to the year the carcass was found as possible (maximum differences between year of death and spatial data were six years for human population and five years for agricultural land use).

2.5. Data analysis

We performed all statistical analyses in R, version 3.2.0 (R Core Team, 2016). We analyzed the data using censored data techniques (R packages NADA; Lee, 2017, and Survival; Therneau, 2018) as some AR concentrations were below the LOD of the LC-MS.

2.5.1. Individual AR and total SGAR concentration

We used a Kaplan-Meier cumulative probability distribution (NADA function 'cenfit') to calculate censored summary statistics (mean, median and standard error) of each AR compound and the total SGAR concentration for each individual eagle. We also calculated analogous standard (non-censored) summary statistics for only the eagles in which ARs were detected. Creating these analogous summary statistics facilitated comparisons among our study and prior work as other studies have used this approach (e.g. Hughes et al., 2013; Lopez-Perea and Mateo, 2018). To facilitate comparisons to other studies, we calculated summary statistics on a wet weight basis. To do this we converted dry weight concentrations (provided in mg/kg dw) to wet weight (mg/kg ww) by multiplying the dry weight concentrations by the dry to wet weight ratios of each sample.

We used total SGAR concentrations to estimate the effects of over-all SGAR contamination due to their similar mode of action (Rattner and Harvey, 2021). FGARs were not included in the summed concentrations due to large differences in molecular weight, potency and half-life compared to SGARs (Rattner and Harvey, 2021). To estimate potential toxicological effects of the total SGAR concentrations detected, we used published contamination thresholds (see Lohr, 2018) as follows: (i) 0.001–0.01 mg/kg ww, probably no toxicity; (ii) 0.01–0.1 mg/kg ww, unlikely lethal/possible toxicity; (iii) 0.1–0.5 mg/kg ww, possibly lethal/likely toxicity; (iv) 0.5–0.7 ww, probably lethal; (v) >0.7 mg/kg ww, lethal. We used the converted wet weight concentrations for this evaluation as the thresholds were based on wet weight concentrations.

2.5.2. Correlates of degree and likelihood of exposure

We explored relationships between the extrinsic and intrinsic explanatory variables (age, sex, year of death, human population density, and proportion of agricultural area) and total SGAR concentration with left-censored regression models (Helsel, 2012; Survival function 'survreg'). We assigned censored data the corresponding LOD value with an indicator variable denoting the observation as below the LOD. Uncensored data were assigned the total liver SGAR concentration measured by the LC-MS and an indicator variable denoting the observation as not censored. The correlation of predictor variables was checked

Table 1

Summary statistics describing liver AR concentrations of each AR assessed and total liver SGAR concentration of Tasmanian wedge-tailed eagle carcasses collected between 1996 and 2018. Non-censored and censored summary statistics are presented. Non-censored statistics were calculated using only the eagles with detected AR concentrations. Censored summary statistics consider all individuals and account for unknown values below the corresponding limit of quantification (LOQ). All summary statistics are provted on a wet weight basis.

before inclusion in the models, and any correlated predictors (Pearson's r > 0.3) were not included in the same model. The dependent variable in these models was the total liver SGAR concentration (mg/kg dw) for each sample. Our initial model set included all possible combinations of submodels. We used corrected Akaike's Information Criterion (AICc) to rank model performance. We excluded models in the candidate set if they had an AICc value greater than six Δ AICc (Richards, 2005). The use of AICc as the sole selection criterion may select overly complex models, thus we considered only those models that had AICc values smaller than all the simpler models within which they were nested (Richards, 2008). We based biological inferences on the coefficients of the top-performing model and considered a parameter to have strong support if it was included in all candidate models.

We also explored the relationship between the same suite of extrinsic and intrinsic predictor variables with the probability of AR residues (both of SGARs and FGARs) being detected using a binomial generalized linear model with logit link function. The dependent variable in these models was whether the eagles were exposed (AR concentrations >LOD) or unexposed (AR concentrations <LOD). We again considered all possible parameter combinations and retained models in the candidate set that were both within six Δ AICc and had AICc values smaller than all the simpler models within which they were nested.

3. Results

We analyzed tissue from 50 eagle carcasses that were collected between 1996 and 2018, although most were collected after 2006 (n = 37). All birds were successfully sexed and aged, with 41 eagles identified as pre-adult, and 22 as female. Data available for the sampled carcasses included location (n = 50; Fig. 1a) and year the carcass was found (n = 50).

3.1. Individual AR and total SGAR concentration

AR residues were detected in 74% of wedge-tailed eagles included in the study (Table 1). Residues of more than one AR compound were detected in 38% of the birds, and 12% of birds had three different compounds detected. The mean total SGAR concentrations of birds in which SGARs were detected was 0.143 mg/kg ww (\pm SE 0.031) and the censored mean of the entire study sample was 0.100 mg/kg ww (\pm SE 0.023). The majority of AR residues were SGARs. Brodifacoum (56% of birds), flocoumafen (40%) and bromadiolone (22%) were the most predominant SGARs detected. FGARs were only detected in three individuals (one of these individuals was also exposed to the SGAR

	Brodifacoum	Bromadiolone	Coumatetralyl	Difenacoum	Difethialone	Flocoumafen	Pindone	Warfarin	Total SGAR
LOQ (mg/kg)	0.005	0.001	0.002	0.0025	0.010	0.0025	0.025	0.002	NA
Birds exposed (%)	28/50 (56%)	11/50 (22%)	1/50 (2%)	0/50 (0%)	0/50 (0%)	20/50 (40%)	0/50 (0%)	2/50 (4%)	37/50 (74%)
Max (mg/kg ww)	0.635	0.241	0.014	0.000	0.000	0.348	0.000	0.002	0.651
Min (mg/kg ww)	0.003 ^a	0.003	0.014 ^a	0.000	0.000	0.002 ^a	0.000	0.001 ^a	0.002
Not censored									
Mean (mg/kg ww)	0.136	0.045	0.014	0.000	0.000	0.035	0.000	0.002	0.143
Median (mg/kg ww)	0.072	0.023	0.014	0.000	0.000	0.003	0.000	0.002	0.074
SE (mg/kg ww)	0.030	0.021	0.000	0.000	0.000	0.022	0.000	0.001	0.031
Censored									
Mean (mg/kg ww)	0.077	0.012	0.014	NA	NA	0.015	NA	0.002	0.100
Median (mg/kg ww)	0.011	<loq< td=""><td><loq< td=""><td><loq< td=""><td><loq< td=""><td><loq< td=""><td><loq< td=""><td><loq< td=""><td>0.017</td></loq<></td></loq<></td></loq<></td></loq<></td></loq<></td></loq<></td></loq<>	<loq< td=""><td><loq< td=""><td><loq< td=""><td><loq< td=""><td><loq< td=""><td><loq< td=""><td>0.017</td></loq<></td></loq<></td></loq<></td></loq<></td></loq<></td></loq<>	<loq< td=""><td><loq< td=""><td><loq< td=""><td><loq< td=""><td><loq< td=""><td>0.017</td></loq<></td></loq<></td></loq<></td></loq<></td></loq<>	<loq< td=""><td><loq< td=""><td><loq< td=""><td><loq< td=""><td>0.017</td></loq<></td></loq<></td></loq<></td></loq<>	<loq< td=""><td><loq< td=""><td><loq< td=""><td>0.017</td></loq<></td></loq<></td></loq<>	<loq< td=""><td><loq< td=""><td>0.017</td></loq<></td></loq<>	<loq< td=""><td>0.017</td></loq<>	0.017
SE (mg/kg ww)	0.019	0.005	NA	NA	NA	0.009	NA	0.000	0.023

^a Trace value.



Fig. 2. Total liver SGAR concentrations (mg/kg ww) for Tasmanian wedge-tailed eagles that died between 1996 and 2018 (n = 50). The number of eagle carcasses with liver SGAR concentrations within each toxicity threshold proposed by Lohr (2018) is presented.

flocoumafen). Warfarin was detected at very low concentrations (<0.01 mg/kg ww) in two birds and coumatetralyl was detected in one bird.

We recorded potentially lethal total liver SGAR concentrations (>0.1 mg/kg ww; Newton et al., 1999) in 11 of the wedge-tailed eagles sampled (22%; 31% of those in which SGARs were detected; Fig. 2). Furthermore, concentrations were above probably lethal levels of >0.5 mg/kg ww in 4% of the eagles (6% of those in which SGARs were detected). That said, liver AR concentrations do not allow the confirmation of lethality without a necropsy to identify signs of toxicity.

3.2. Correlates of degree of exposure

The top-performing censored regression model suggested that total liver SGAR concentration (mg/kg dw) was driven most strongly by the year the carcass was found, the amount of agricultural area, and the human population density in the area around where the carcass was found (see Appendix Table G.1). This model was 42.83 times more likely than the null model. A simpler model that excluded human population density was also retained in the candidate model set (Table 2). The year the carcass was found and agricultural area were included in both candidate models, suggesting that these variables were the most important to explaining total liver AR concentration. Coefficients of the best performing model indicated that year of death, agricultural area, and human population density were all positively associated with total AR

Table 2

Candidate models relating total liver SGAR concentrations and probability of ARs being detected in Tasmanian wedge-tailed eagle carcasses collected between 1996 and 2018 to intrinsic and extrinsic factors.^a

Rank	Model variables	df	AICc	AICc weight		
Total liver AR concentration						
1	Year of death (+); Agricultural area (+); Human population density (+)	5	93.170	0.795		
2	Year of death (+); Agricultural area (+)	4	95.885	0.205		
Probability of exposure						
1	Year of death $(+)$; Agricultural area $(+)$	3	55.010	0.712		
2	Year of death (+)	2	57.471	0.208		
3	Null model	1	59.389	0.079		

^a Only models that were both within six Δ AlCc and had AlCc values smaller than all the simpler models within which they were nested were retained in the candidate model set (Richards, 2008). For year of death, "+" indicates association to carcasses found more recently.

concentration (Table 3, Fig. 3a). The model suggested that a 10% increase in agricultural habitat in the area around where the bird died would result in an increase in liver AR concentrations by a factor of 2.11. Likewise, each later year in the study was estimated to increase AR concentrations by a factor of 1.40. The relationship between total AR concentration and human population density suggested an increase in 100 habitants per km² would increase total AR concentration by a factor of 7.23.

3.3. Correlates of likelihood of exposure

The top-performing binomial model to explain the probability of an eagle being exposed to ARs included the year the carcass was found and the proportion of agricultural area within the area around where the carcass was found (see Appendix Table G.2). The candidate model set included two simpler models, including the null model (Table 2), al-though the top-performing model was 8.9 times more likely than the null model based on AICc weight. Coefficients of the top-performing model indicated that the probability of ARs being detected increased with carcasses found more recently and in areas with higher proportions of agricultural area (Table 3). The odds of ARs being detected in a carcass were 1.46 times greater for each 10% increase in agricultural habitat proportion in the area around where the bird died and 1.21 times greater for each advancing year of the study period (Fig. 3b, Appendix Fig. G.1).

4. Discussion

The frequency and magnitude of AR exposure in Tasmanian wedgetailed eagles, and their correlation to agricultural areas and human population density, have several implications for our understanding of rodenticide exposure and the Tasmanian ecosystem. First, rodenticide exposure is high among these birds, suggesting that rodenticides are frequently finding their way into top predators in the ecosystem. Furthermore, extrinsic (i.e. agricultural area, human population density, and year of death) rather than intrinsic factors (i.e. age, sex) influence the probability of exposure to ARs and total SGAR concentration. These findings illustrate how AR exposure of the Tasmanian wedgetailed eagle is driven by anthropogenic processes and thus identify directions to solve this conservation problem.

4.1. Individual AR and total SGAR concentration

The high prevalence of SGARs detected in our study is consistent with research implicating SGARs as the predominant cause of non-

Table 3

Model coefficients for top performing models describing the estimated effect of each variable on total liver SGAR concentrations (censored regression) and the probability of an eagle being exposed to ARs (binomial probability).

Model	Parameter	Estimate	95% CI lower	95% CI upper	Ζ
Total liver AR concentration					
Probabi	Intercept Year of death Agricultural area Human population density lity of exposure	-12.105 0.337 0.749 1.978	-16.669 0.137 0.248 0.306	-7.542 0.536 1.249 3.650	-5.20 3.31 2.93 2.32
	Intercept Year of death Agricultural area	—2.945 0.193 0.375	-6.313 0.053 0.035	—0.165 0.365 0.781	-1.93 2.48 2.02

target AR exposure of predators (Lohr, 2018; López-Perea et al., 2015). SGARs brodifacoum, bromadiolone, and flocoumafen accounted for 99.6% of the total AR concentrations observed in the Tasmanian wedge-tailed eagle. The first two of these are the AR compounds most commonly identified in non-target predators in numerous ecosystems worldwide (Hosea, 2000; Koivisto et al., 2016; Langford et al., 2013; Ruiz-Suárez et al., 2014; Sharp et al., 2005). The extent of the flocoumafen contamination we detected is more surprising and represents one of the highest exposure rates documented globally (see Appendix Table G.3). Flocoumafen is only available through wholesale outlets in Tasmania, suggesting that agricultural asset protection and



Fig. 3. a) Predicted response of total liver SGAR concentrations (mg/kg dw) in Tasmanian wedge-tailed eagle carcasses as a function of the proportion of agricultural land area and mean human population density in the area around where the bird died. The three lines are the estimated response of liver AR concentration with human population per km² held at three levels. Year of carcass discovery is held at its mean. b) Logistic plot of the effect of year of death on the probability of AR exposure. Agricultural land area is held at its mean.

professional pest controllers could be important sources of non-target AR exposure in Australia.

The low concentrations and frequency of detections of FGARs also corresponds with findings for other species, both in Australia and globally (Cypher et al., 2014; Lohr, 2018; Murray, 2020, 2017). This was of particular interest in the case of the FGAR pindone. We expected pindone to be the AR most frequently detected in wedge-tailed eagles, since it targets a common prey item for the species (the European rabbit). Our low detection of FGARs could be due to their characteristics - their shorter half-life and lower toxicity – and, in the case of pindone, its localized use in targeted control efforts (Lohr, 2018). Although this low rate of detection may suggest FGARs pose a lower risk of non-target exposure, their characteristics may also impede their detection in studies using opportunistic sampling and prolonged tissue storage (Herring et al., 2017; Rattner et al., 2014), consequently underestimating their true prevalence in the Australian environment.

The frequency and magnitude of AR exposure in the Tasmanian wedge-tailed eagle was high for an Aquila species. Raptor studies showing comparable AR detection rates typically involve smaller species known to be at risk due to their dietary specialization on rodents (Christensen et al., 2012; López-Perea et al., 2015; Walker et al., 2011). The proportion of birds we observed with concentrations >0.2 mg/kg ww (16%) is substantially higher than that found in congeners (0-6%; Hosea, 2000; Langford et al., 2013; Sánchez-Barbudo et al., 2012), and the highest concentration of an SGAR we detected in an individual (0.635 mg/kg ww of brodifacoum) is substantially higher than the highest concentration of an AR previously reported in an Aquila species (0.154 mg/kg ww of bromadiolone; Langford et al., 2013). Both the censored mean SGAR concentrations of all eagles sampled (0.100 mg/kg ww) and the mean only of those with detected SGAR levels (0.143 mg/kg ww) were higher than mean concentrations reported for congeners (0.006–0.073 mg/kg ww; Langford et al., 2013; Sánchez-Barbudo et al., 2012), but lie within the range of values reported for other raptors exposed to SGARs (0.005-0.413 mg/kg ww; Thomas et al., 2011; Christensen et al., 2012; Lohr, 2018).

The high exposure to SGARs in the Tasmanian wedge-tailed eagle, a species not known to regularly prey upon synanthropic rodents, supports the suggestion that apex predators are vulnerable to SGARs (López-Perea et al., 2015; Riley et al., 2007). The long halflife and persistence of SGARs gives these compounds the capacity to move through food chains (López-Perea et al., 2015), a theory evidenced by the presence of ARs in apex predators (Riley et al., 2007). The wedge-tailed eagle preys upon several carnivorous species that are known to consume synanthropic rodents. For example predatory and scavenging species such as forest ravens (Corvus tasmanicus), kookaburras (Dacelo novaeguineae), common brush-tail possums (Trichosurus vulpecula), cats (Felis catus), and other raptors have been recorded in wedge-tailed eagle diets (Marchant and Higgins, 1993). The potential for SGARs to move through multiple trophic levels may therefore be causing extensive contamination of Tasmania's terrestrial food chains (Thomas et al., 2011). If this is the case, then numerous other predatory species may be at risk in the region, including the endangered Tasmanian devil and eastern quoll (Dasyurus viverrinus; IUCN, 2020).

The high exposure we detected may also be driven by the improper use of ARs and non-target AR vectors. The use of SGARs in Australia does not require a license, products can be easily purchased in large quantities, and awareness of use guidelines may be low. If SGARs are not being used as directed, numerous non-rodent species may consume the poisons and act as AR vectors to predators. Furthermore, ARs have recently been detected in Australian reptile species; this exposure could be through direct consumption of ARs used correctly, since these species are small enough to enter AR bait boxes (Lettoof et al., 2020). Reptiles are prey for wedge-tailed eagles, as well as other predators, and may therefore have a role as AR vectors in Australia.

4.2. Correlates of AR exposure

The positive association between hepatic AR concentrations and human population density and agricultural land use may indicate localized use that is having wider scale effects. AR residues in predators have been linked to human population density (López-Perea et al., 2018, 2015), the amount of urbanized area (Coeurdassier et al., 2019; Lohr, 2018; Serieys et al., 2015), and the amount of both arable and pastoral agriculture (Coeurdassier et al., 2019; López-Perea et al., 2018; Sainsbury et al., 2018). These relationships are unsurprising in study species known to use urban and agricultural habitats. However, Tasmanian wedge-tailed eagles are less associated with densely populated areas. Although human population growth has been relatively low in Tasmania for the past two decades, there has been an increase in the number of residences built in more rural and natural areas and agricultural development has expanded (Australian Bureau of Statistics, 2018). Such practices may introduce ARs into more natural areas. Furthermore, if ARs are passing through multiple trophic levels, they will spread more widely from the initial bait. The effects of these more remote developments and agricultural activities may therefore have incommensurately greater effects on predatory species than suggested by the landscape footprint.

4.3. Recent increases in AR exposure

The higher total SGAR concentrations and probability of AR exposure of the birds that had died more recently could be due to either the increased exposure to ARs over time or the degradation of the compounds with prolonged storage. Although SGAR residues are stable within tissues over the short-term (Gallocchio et al., 2014; Jin and Chen, 2006), the effects of long-term -20 °C freezer storage on tissue residues is not well understood, with studies documenting various rates of degradation (e.g. 6-41% over 0.5-3 years; P. Fisher unpublished data; Vindenes et al., 2008). Despite this, patterns in the AR concentrations we detected are consistent with increased probability of exposure over time. There would need to be a substantial reduction in AR residues (much greater than the degradation rates documented) for an ARexposed bird to be considered unexposed in our study, as the lowest AR concentration we detected was still 200% greater than the associated LOD. Consequently, the increased AR concentrations in Tasmanian wedge-tailed eagles that had died more recently is more likely due to increases or changes in AR use in Tasmania than to sample degradation. However, there is no information available on the volume of ARs used in Australia (Lohr and Davis, 2018), which impedes our quantification of the relationship between AR application and non-target AR exposure.

4.4. Conservation implications

Our results indicate that AR exposure is likely a significant factor to consider in the conservation management for the Tasmanian wedgetailed eagle. This is true even given the potential biases inherent to the non-random carcass collection we relied on to gather samples. AR studies using opportunistic samples may inflate the proportion of animals with sub-lethal AR concentrations detected and underestimate the proportion of birds detected with fatal levels (Lohr, 2018; Newton et al., 1990). We found exposure at rates that are high compared to other studies using similar sampling methods.

The use of AR concentration thresholds to interpret the likely physiological result has limitations due to inter- and intra-specific variation in susceptibility to toxicity (Rattner and Harvey, 2021; Thomas et al., 2011). That said, concentrations in 22% of the birds we studied were well above the potentially lethal range reported for European barn owls (*Tyto alba*; >0.1 mg/kg; Newton et al., 1999). Furthermore, 56% had levels that can cause symptoms of toxicity in other species (>0.01 mg/kg; Lohr, 2018; Murray, 2018);. These comparative data therefore suggest that the level of exposure we detected indicates that AR exposure could be influential to survival and possibly conservation of these birds.

Our findings underscore the importance of efforts to address nontarget AR exposure in Australian wildlife. SGARs are currently registered for domestic (non-professional) use in Australia (Australian Pesticides and Veterinary Medicines Authority, 2019), despite increasing regulation and monitoring in other countries (USEPA, 2008). Increased legislative control of SGARs and removal from public retail have been suggested as steps to reduce the ecological effects of SGAR use in Australia (Lohr and Davis, 2018). However, our findings of an association between agriculture and AR concentrations in the Tasmanian wedge-tailed eagle, as well as widespread contamination of an AR not readily available for residential use (flocoumafen), suggests that professional pest control may also be an important cause of non-target AR exposure. Addressing mechanisms of spread from both professional and non-professional application of SGARs may therefore be important to reducing AR exposure in Tasmanian wedge-tailed eagles and other Australian wildlife.

Disclaimer

Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the US Government.

Data accessibility

Data available as electronic supplementary material.

Funding

This work was supported by Woolnorth Wind Farms, TasNetworks, New Forests, Norske Skog, and the Holsworth Wildlife Research Endowment.

CRediT authorship contribution statement

James M. Pay: Conceptualization, Methodology, Formal analysis, Investigation, Data curation, Visualization, Writing – original draft, Project administration, Funding acquisition, Writing – review & editing. Todd E. Katzner: Conceptualization, Methodology, Investigation, Writing – original draft, Supervision, Formal analysis, Writing – review & editing. Clare E. Hawkins: Conceptualization, Writing – original draft, Supervision. Leon A. Barmuta: Formal analysis, Writing – review & editing. William E. Brown: Investigation, Resources, Writing – review & editing. Jason M. Wiersma: Conceptualization, Investigation, Writing – review & editing. Jason M. Wiersma: Conceptualization, Investigation, Writing – review & editing. Amelia J. Koch: Conceptualization, Writing – original draft, Supervision, Funding acquisition. Nick J. Mooney: Investigation, Resources, Writing – review & editing. Conceptualization, Writing – original draft, Supervision, Funding acquisition. Nick J. Mooney: Investigation, Resources, Writing – original draft, Supervision, Project administration, Funding acquisition.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This study was possible due to the long-term storage of Tasmanian wedge-tailed eagle carcasses provided by the Tasmanian Museum and Art Gallery and the Department of Primary Industries, Parks, Water and Environment. We thank the numerous people and organizations that were involved in collecting these carcasses. B. Bauer, J. Harris, and N. Dannemiller assisted necropsies. We also thank Mike Lohr, Francesco Busetti, and Kirstin Proft for their contribution to this

research. Laboratory analyses were carried out by Edith Cowan University Analytical Facility.

Supplementary materials

Supplementary materials to this article can be found online at https://doi.org/10.1016/j.scitotenv.2021.147673.

References

- Australian Bureau of Statistics, 2018. Regional Population Growth, Australia 2016 (3218). (Canberra).
- Australian Pesticides and Veterinary Medicines Authority, 2019. Public chemical registration information system search. (Canberra).
- Bell, P., Mooney, N., 1998. Wedge-Tailed Eagle Recovery Plan 1998–2003. Department of Primary Industries, Parks, Water and Environment, Hobart.
- Birks, J.D.S., 1998. Secondary rodenticide poisoning risk arising from winter farmyard use by the European polecat *Mustela putorius*. Biol. Conserv. 85, 233–240. https://doi.org/ 10.1016/S0006-3207(97)00175-4.
- Brakes, C.R., Smith, R.H., 2005. Exposure of non-target small mammals to rodenticides: short-term effects, recovery and implications for secondary poisoning. J. Appl. Ecol. 42, 118–128. https://doi.org/10.1111/j.1365-2664.2005.00997.x.
- Christensen, T.K., Lassen, P., Elmeros, M., 2012. High exposure rates of anticoagulant rodenticides in predatory bird species in intensively managed landscapes in Denmark. Arch. Environ. Contam. Toxicol. 63, 437–444. https://doi.org/10.1007/ s00244-012-9771-6.
- Coeurdassier, M., Villers, A., Augiron, S., Sage, M., Couzi, F.X., Lattard, V., Fourel, I., 2019. Pesticides threaten an endemic raptor in an overseas French territory. Biol. Conserv. 234, 37–44. https://doi.org/10.1016/j.biocon.2019.03.022.
- Commonwealth of Australia, 1999. Environment Protection and Biodiversity Conservation Act 1999 - EPBC Act List of Threatened Fauna (Canberra).
- Cypher, B.L., McMillin, S.C., Westall, T.L., Van Horn Job, C., Hosea, R.C., Finlayson, B.J., Kelly, E.C., 2014. Rodenticide exposure among endangered kit foxes relative to habitat use in an urban landscape. Cities Environ. 7, 1–21.
- Debus, S.J.S., Hatfield, T.S., Ley, A.J., Rose, A.B., 2007. Breeding biology and diet of the wedge-tailed eagle *Aquila audax* in the New England region of New South Wales. Aust. F. Ornithol. 24, 93–120.
- DPIPWE, 2015. Tasmania land use 2001-2015 [WWW document]. NRM Data Libr. URL https://listdata.thelist.tas.gov.au/opendata/ (accessed 2.15.19).
- Erickson, W., Urban, D., 2004. Potential Risk of Nine Rodenticides to Birds and Mammals: A Comparative Approach. Office of Pesticides Programs Environmental Fate and Effects Division, Washington, D.C.
- Fourel, I., Sage, M., Benoit, E., Lattard, V., 2018. Liver and fecal samples suggest differential exposure of red fox (*Vulpes vulpes*) to trans- and cis-bromadiolone in areas from France treated with plant protection products. Sci. Total Environ. 622–623, 924–929. https://doi.org/10.1016/j.scitotenv.2017.12.053.
- Gallocchio, F., Basilicata, L., Benetti, C., Angeletti, R., Binato, G., 2014. Multi-residue determination of eleven anticoagulant rodenticides by high-performance liquid chromatography with diode array/fluorimetric detection: investigation of suspected animal poisoning in the period 2012-2013 in north-eastern Italy. Forensic Sci. Int. 244, 63–69. https://doi.org/10.1016/j.forsciint.2014.08.012.
- Helsel, D.R., 2012. Statistics for Censored Environmental Data Using Minitab and R. Wiley, Hoboken.
- Herring, G., Eagles-Smith, C.A., Buck, J., 2017. Characterizing golden eagle risk to lead and anticoagulant rodenticide exposure: a review. J. Raptor Res. 51, 273–292. https://doi. org/10.3356/JRR-16-19.1.
- Hindmarch, S., Elliott, J.E., 2018. Ecological factors driving uptake of anticoagulant rodenticides in predators. In: van den Brink, N.W., Elliot, J.E., Shore, R.F., Rattner, B.A. (Eds.), Anticoagulant Rodenticides and Wildlife. Springer International, Cham, pp. 229–258.
- Hindmarch, S., Elliott, J.E., Mccann, S., Levesque, P., 2017. Habitat use by barn owls across a rural to urban gradient and an assessment of stressors including, habitat loss, rodenticide exposure and road mortality. Landsc. Urban Plan. 164, 132–143. https://doi. org/10.1016/j.landurbplan.2017.04.003.
- Horak, K.E., Fisher, P.M., Hopkins, B., 2018. Pharmacokinetics of anticoagulant rodenticides in target and non-target organisms. In: van den Brink, N.W., Elliott, J.E., Shore, R.F., Rattner, B.A. (Eds.), Anticoagulant Rodenticides and Wildlife. Springer International, Cham, pp. 87–108.
- Hosea, R.C., 2000. Exposure of non-target wildlife to anticoagulant rodenticides in California. In: Salmon, T.P., Crabb, A.C. (Eds.), Proceedings of the 19th Vertebrate Pest Conference. University of California, Davis, pp. 236–244.
- Hughes, J., Sharp, E., Taylor, M.J., Melton, L., Hartley, G., 2013. Monitoring agricultural rodenticide use and secondary exposure of raptors in Scotland. Ecotoxicology 22, 974–984. https://doi.org/10.1007/s10646-013-1074-9.
- IUCN, 2020. The IUCN Red List of Threatened Species. Version 2020-2 [WWW Document]. URL www.iucnredlist.org. (Accessed 8.16.20).
- Jacquot, M., Coeurdassier, M., Couval, G., Renaude, R., Pleydell, D., Truchetet, D., Raoul, F., Giraudoux, P., 2013. Using long-term monitoring of red fox populations to assess changes in rodent control practices. J. Appl. Ecol. 50, 1406–1414. https://doi.org/ 10.1111/1365-2664.12151.
- Jin, M., Chen, X., 2006. Rapid determination of three anticoagulant rodenticides in whole blood by liquid chromatography coupled with electrospray ionization mass spectrometry. Rapid Commun. Mass Spectrom. 20, 2741–2746. https:// doi.org/10.1002/rcm.

- Koivisto, E., Koivisto, P., Hanski, I.K., Korkolainen, T., Vuorisalo, T., Karhilahti, A., Loivamaa, I., Koivisto, S., 2016. Prevalence of anticoagulant rodenticides in non-target predators and scavengers in Finland. Report of the Finnish Safety and Chemicals Agency (Tukes) (Helsinki).
- Langford, K.H., Reid, M., Thomas, K.V., 2013. The occurrence of second generation anticoagulant rodenticides in non-target raptor species in Norway. Sci. Total Environ. 450–451, 205–208. https://doi.org/10.1016/j.scitotenv.2013.01.100.
- Lee, L., 2017. NADA: Nondetects and Data Analysis for Environmental Data. R package version 1.6-1.
- Lee, M., Morfini, M., Negrier, C., Chamouard, V., 2006. The pharmacokinetics of coagulation factors. Haemophilia 12 Suppl, 1–7.
- Lettoof, D.C., Lohr, M.T., Busetti, F., Bateman, P.W., Davis, R.A., 2020. Toxic time bombs: frequent detection of anticoagulant rodenticides in urban reptiles at multiple trophic levels. Sci. Total Environ. 724, 138218. https://doi.org/10.1016/j.scitotenv.2020.138218.
- Lohr, M.T., 2018. Anticoagulant rodenticide exposure in an Australian predatory bird increases with proximity to developed habitat. Sci. Total Environ. 643, 134–144. https://doi.org/10.1016/j.scitotenv.2018.06.207.
- Lohr, M.T., Davis, R.A., 2018. Anticoagulant rodenticide use, non-target impacts and regulation: a case study from Australia. Sci. Total Environ. 634, 1372–1384. https://doi. org/10.1016/j.scitotenv.2018.04.069.
- Lopez-Perea, J.J., Mateo, R., 2018. Secondary exposure to anticoagulant rodenticides and effects on predators. In: van den Brink, N.W., Elliott, J.E., Shore, R.F., Rattner, B.A. (Eds.), Anticoagulant Rodenticides and Wildlife. Springer, Cham, pp. 159–193.
- López-Perea, J.J., Camarero, P.R., Molina-López, R.A., Parpal, L., Obón, E., Solá, J., Mateo, R., 2015. Interspecific and geographical differences in anticoagulant rodenticide residues of predatory wildlife from the Mediterranean region of Spain. Sci. Total Environ. 511, 259–267. https://doi.org/10.1016/j.scitotenv.2014.12.042.
- López-Perea, J.J., Camarero, P.R., Sánchez-Barbudo, I.S., Mateo, R., 2018. Urbanization and cattle density are determinants in the exposure to anticoagulant rodenticides of non-target wildlife. Environ. Pollut. 244, 801–808. https://doi.org/10.1016/I.ENVPOL2018.10.101.
- Marchant, S., Higgins, P.J., 1993. Handbook of Australian, New Zealand and Antarctic Birds. Volume Two—Raptors to Lapwings. Oxford University Press, Melbourne.
- Martin, G.R., Kirkpatrick, W., King, D., Robertson, I., Hood, P., Sutherland, J., 1994. Assessment of the potential toxicity of an anticoagulant, pindone (2-Pivalyl-1,3-Indandione), to some Australian birds. Wildl. Res. 21, 85–93. https://doi.org/10.1071/WR9940085.
- Mcdonald, R.A., Harris, S., Turnbull, G., Brown, P., Fletcher, M., 1998. Anticoagulant rodenticides in stoats (*Mustela erminea*) and weasels (*Mustela nivalis*) in England. Environ. Pollut, 103, 17–23.
- Mooney, N., 2017. Risks of anticoagulant rodenticides to Tasmanian raptors. Tasmanian Bird Rep. 38, 17–25.
- Mooney, N., Holdsworth, M., 1991. The effects of disturbance on nesting wedge-tailed eagles (Aquila audax fleayi) in Tasmania. Tasforests 3, 15–31.
- Murray, M., 2017. Anticoagulant rodenticide exposure and toxicosis in four species of birds of prey in Massachusetts, USA, 2012–2016, in relation to use of rodenticides by pest management professionals. Ecotoxicology 26, 1041–1050. https://doi.org/ 10.1007/s10646-017-1832-1.
- Murray, M., 2018. Ante-mortem and post-mortem signs of anticoagulant rodenticide toxicosis in birds of prey. In: van den Brink, N., Elliott, J.E., Shore, R.F., Rattner, B.A. (Eds.), Anticoagulant Rodenticides and Wildlife. Springer International, Cham, pp. 109–134.
- Murray, M., 2020. Continued anticoagulant rodenticide exposure of red-tailed hawks (*Buteo jamaicensis*) in the northeastern United States with an evaluation of serum for biomonitoring. Environ. Toxicol. Chem. 39, 2325–2335. https://doi.org/10.1002/etc.4853.
- Newton, I., Wyllie, I., Freestone, P., 1990. Rodenticides in British barn owls. Environ. Pollut. 68, 101–117.
- Newton, I., Shore, R.F., Wyllie, I., Briks, J.D.S., Dale, L., 1999. Empirical evidence of sideeffects of rodenticides on some predatory birds and mammals. In: Cowan, D.P., Frear, C.J. (Eds.), Advances in Vertebrate Pest Management. Filander-Verlag, Fürth, pp. 347–367.
- Nogeire, T.M., Lawler, J.J., Schumaker, N.H., Cypher, B.L., Phillips, S.E., 2015. Land use as a driver of patterns of rodenticide exposure in modeled kit fox populations. PLoS One 10, e0133351. https://doi.org/10.1371/journal.pone.0133351.
- Pay, J.M., Katzner, T.E., Hawkins, C.E., Koch, A.J., Wiersma, J.M., Brown, W.E., Mooney, N.J., Cameron, E.Z., 2020. High frequency of lead exposure in the population of an endangered top predator, the Tasmanian wedge-tailed eagle (*Aquila audax fleayi*). Environ. Toxicol. Chem. 40, 219–230.
- R Core Team, 2016. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna https://doi.org/10.1007/978-3-540-74686-7.
- Rattner, B.A., Harvey, J.J., 2021. Challenges in the interpretation of anticoagulant rodenticide residues and toxicity in predatory and scavenging birds. Pest Manag. Sci. 77, 604–610. https://doi.org/10.1002/ps.6137.
- Rattner, B.A., Lazarus, R.S., Elliott, J.E., Shore, R.F., van den Brink, N., 2014. Adverse outcome pathway and risks of anticoagulant rodenticides to predatory wildlife. Environ. Sci. Technol. 48, 8433–8445. https://doi.org/10.1021/es501740n.
- Rial-Berriel, C., Acosta-Dacal, A., Cabrera Pérez, M.Á., Suárez-Pérez, A., Melián Melián, A., Zumbado, M., Henríquez Hernández, L.A., Ruiz-Suárez, N., Rodriguez Hernández, Á., Boada, L.D., Macías Montes, A., Luzardo, O.P., 2021. Intensive livestock farming as a major determinant of the exposure to anticoagulant rodenticides in raptors of the Canary Islands (Spain). Sci. Total Environ. 768. https://doi. org/10.1016/j.scitotenv.2020.144386.
- Richards, S.A., 2005. Testing ecological theory using the information-theoretic approach: examples and cautionary results. Ecology 86, 2805–2814. https://doi.org/10.1890/ 05-0074.
- Richards, S.A., 2008. Dealing with overdispersed count data in applied ecology. J. Appl. Ecol. 45, 218–227. https://doi.org/10.1111/j.1365-2664.2007.01377.x.

- Riley, S.P.D., Bromley, C., Poppenga, R.H., Uzal, F.A., Whited, L., Sauvajot, R.M., 2007. Anticoagulant exposure and notoedric mange in bobcats and mountain lions in urban Southern California. J. Wildl. Manag. 71, 1874–1884. https://doi.org/ 10.2193/2005-615.
- Ruiz-Suárez, N., Henríquez-Hernández, L.A., Valerón, P.F., Boada, L.D., Zumbado, M., Camacho, M., Almeida-González, M., Luzardo, O.P., 2014. Assessment of anticoagulant rodenticide exposure in six raptor species from the Canary Islands (Spain). Sci. Total Environ. 485–486, 371–376. https://doi.org/10.1016/j.scitotenv.2014.03.094.
- Ruiz-Suárez, Melero, Giela, Y., Henríquez-Hernández, A., Sharp, L.A., Boada, E., Taylor, L.D., Camacho, M.J., Lambin, M., Luzardo, X., Hartley, O.P., 2016. Rate of exposure of a sentinel species, invasive American mink (*Neovison vison*) in Scotland, to anticoagulant rodenticides. Sci. Total Environ. 569–570, 1013–1021. https://doi.org/10.1016/j. scitotenv.2016.06.109.
- Sainsbury, K.A., Shore, R.F., Schofield, H., Croose, E., Hantke, G., Pereira, M.G., Sleep, D., Kitchener, A.C., McDonald, R.A., 2018. Long-term increase in secondary exposure to anticoagulant rodenticides in European polecats *Mustela putorius* in Great Britain. Environ. Pollut. 236, 689–698. https://doi.org/10.1016/j.envpol.2018.02.004.Sánchez-Barbudo, I.S., Camarero, P.R., Mateo, R., 2012. Primary and secondary poisoning
- Sánchez-Barbudo, I.S., Camarero, P.R., Mateo, R., 2012. Primary and secondary poisoning by anticoagulant rodenticides of non-target animals in Spain. Sci. Total Environ. 420, 280–288. https://doi.org/10.1016/j.scitotenv.2012.01.028.
- Serieys, L.E.K., Armenta, T.C., Moriarty, J.G., Boydston, E.E., Lyren, L.M., Poppenga, R.H., Crooks, K.R., Wayne, R.K., Riley, S.P.D., 2015. Anticoagulant rodenticides in urban bobcats: exposure, risk factors and potential effects based on a 16-year study. Ecotoxicology 24, 844–862. https://doi.org/10.1007/s10646-015-1429-5.
- Sharp, E., T, A.G., Hunter, K., 2005. The Environmental Impact of Anticoagulant Rodenticide Use on Wildlife in Scotland. Scottish Agricultural Science Agency, Edinburgh.
- Shore, R.F., Birks, J.D.S., Afsar, A., Wienburg, C.L., Kitchener, A.C., 2003. Spatial and temporal analysis of second-generation anticoagulant rodenticide residues in polecats

(*Mustela putorius*) from throughout their range in Britain, 1992–1999. Environ. Pollut. 122, 183–193. https://doi.org/10.1016/S0269-7491(02)00297-X.

- State Government of Tasmania, 1995. Threatened Species Protection Act 1995 Schedule 3 Taxa of Flora and Fauna which Are Endangered. State Government of Tasmania, Hobart.
- Therneau, T.M., 2018. survival: Survival Analysis. R package version 2.43-3.
- Thomas, P.J., Mineau, P., Shore, R.F., Champoux, L., Martin, P.A., Wilson, L.K., Fitzgerald, G., Elliott, J.E., 2011. Second generation anticoagulant rodenticides in predatory birds: probabilistic characterisation of toxic liver concentrations and implications for predatory bird populations in Canada. Environ. Int. 37, 914–920. https://doi.org/10.1016/j. envint.2011.03.010.
- Thomas, P.J., Eccles, K.M., Mundy, L.J., 2017. Spatial modelling of non-target exposure to anticoagulant rodenticides can inform mitigation options in two boreal predators inhabiting areas with intensive oil and gas development. Biol. Conserv. 212, 111–119. https://doi.org/10.1016/i.biocon.2017.06.005.
- 111–119. https://doi.org/10.1016/j.biocon.2017.06.005. Threatened Species Section, 2006. Threatened Tasmanian Eagles Recovery Plan: 2006–2010. Department of Primary Industries, Parks, Water and Environment, Hobart.
- USEPA, 2008. Risk Mitigation Decision for Ten Rodenticides. USEPA Office of Prevention, Pesticides and Toxic Substances, Washington, D.C.
- van den Brink, N., Elliott, J.E., Shore, R.F., Rattner, B.A., 2018. Anticoagulant Rodenticides and Wildlife. Springer International, Cham https://doi.org/10.1007/978-3-319-64377-9.
- Vindenes, V., Karinen, R., Hasvold, I., Bernard, J.P., Mørland, J.G., Christophersen, A.S., 2008. Bromadiolone poisoning: LC-MS method and pharmacokinetic data. J. Forensic Sci. 53, 993–996. https://doi.org/10.1111/j.1556-4029.2008.00737.x.
- Walker, L.A., Chaplow, J.S., Llewellyn, N.R., Pereira, M.G., Potter, E.D., Sainsbury, A.W., Shore, R.F., 2011. Anticoagulant Rodenticides in Predatory Birds 2011: A Predatory Bird Monitoring Scheme (PBMS) Report. Centre for Ecology & Hydrology, Lancaster.